

Assessment of Foveal Avascular Zone Size in Diabetic Retinopathy Using Optical Coherence Tomography Angiography (OCTA)

¹Dr. Muhammad Usman, ²Dr. Faisal Nawaz, ³Dr. Muhammad Zaheer ullah Babar, ⁴Dr. Asif Ali, ⁵Mr. Muhammad Harris Tariq

¹Assistant professor. Department of Ophthalmology, Peshawar Medical College, Riphah International University

²Assistant professor, Department of Ophthalmology, Peshawar Medical College, Riphah International University

³Senior Registrar, Department of Ophthalmology, Peshawar Medical College, Riphah International University

⁴Assistant professor, Department of Ophthalmology, Peshawar Medical College, Riphah International University

⁵Optometrist, Department of Ophthalmology. Peshawar Medical College. Riphah International University

ABSTRACT:

Background: Diabetic retinopathy (DR) is a leading cause of vision impairment, characterized by microvascular changes in the retina. The foveal avascular zone (FAZ) plays a critical role in maintaining central vision, and its enlargement is associated with disease severity in DR. Optical coherence tomography angiography (OCTA) is a non-invasive imaging modality that allows for detailed evaluation of retinal microvasculature, including the FAZ.

Aim: This study aimed to assess the size of the foveal avascular zone in patients with different stages of diabetic retinopathy using OCTA and to determine its correlation with disease progression.

Methods: A cross-sectional study was conducted on 30 patients diagnosed with various stages of diabetic retinopathy. OCTA was used to measure the FAZ area in both the superficial and deep capillary plexus. Patients were classified into non-proliferative DR (mild, moderate, and severe) and proliferative DR groups. Statistical analyses were performed to evaluate differences in FAZ size across the groups and to assess the correlation between FAZ enlargement and DR severity.

Results: FAZ size was found to be significantly larger in patients with advanced stages of diabetic retinopathy. The average FAZ area in the superficial plexus increased progressively from mild non-proliferative DR ($0.26 \pm 0.07 \text{ mm}^2$) to proliferative DR ($0.42 \pm 0.11 \text{ mm}^2$). Similarly, the deep capillary plexus showed a significant increase in FAZ size as DR severity advanced. A strong positive correlation ($r = 0.81, p < 0.001$) was observed between FAZ enlargement and DR progression.

Conclusion: OCTA provided valuable insights into the microvascular changes occurring in diabetic retinopathy. The enlargement of the foveal avascular zone was closely associated with disease severity, making it a potential biomarker for monitoring DR progression. Early detection of FAZ changes using OCTA could facilitate timely intervention and prevent further visual deterioration.

Keywords: Diabetic retinopathy, foveal avascular zone, optical coherence tomography angiography, retinal microvasculature, disease progression, OCTA

INTRODUCTION:

Diabetic retinopathy (DR) has long been recognized as a leading cause of visual impairment and blindness in individuals with diabetes. The increasing global prevalence of diabetes has amplified the burden of diabetic retinopathy, making it a critical public health concern. Previous studies have identified microvascular changes, such as capillary non-perfusion and increased vascular permeability, as key contributors to the progression of diabetic retinopathy [1]. Traditionally, these microvascular changes were assessed using techniques like fundus fluorescein angiography (FFA), which provided valuable

insights but had limitations, including invasiveness and potential complications. The development of optical coherence tomography angiography (OCTA), a non-invasive imaging modality, provided a more detailed and safer alternative for assessing retinal vascular structures, particularly the foveal avascular zone (FAZ), an important biomarker in diabetic retinopathy [2].

The foveal avascular zone, a capillary-free area located at the center of the macula, is vital for maintaining high visual acuity. In diabetic retinopathy, alterations in FAZ size and morphology have been linked to disease progression and visual impairment [3]. Several studies have demonstrated that the enlargement of the FAZ is correlated with the severity of diabetic retinopathy, making it a potential marker for early detection and monitoring of the disease. Optical coherence tomography angiography (OCTA) enabled researchers to measure FAZ size and assess retinal microvascular changes with high precision, offering a significant advantage over conventional imaging methods [4].

Optical coherence tomography angiography (OCTA) employed motion contrast imaging to capture dynamic blood flow within the retinal and choroidal vasculature. Its ability to visualize both superficial and deep capillary plexuses without the need for dye injection reduced the risks associated with FFA and provided detailed information on the retinal microvasculature [5]. In diabetic retinopathy, OCTA enabled the assessment of capillary dropout, microaneurysms, and FAZ alterations, all of which contributed to a comprehensive understanding of the disease's pathophysiology.

Previous studies had explored the relationship between FAZ size and diabetic retinopathy stages, but the findings were varied. Some researchers reported a progressive increase in FAZ size with worsening diabetic retinopathy, while others suggested that FAZ alterations might be detectable even in early stages of the disease [6]. Additionally, the impact of diabetic macular edema (DME), another complication of diabetic retinopathy, on FAZ morphology had been investigated. DME, characterized by fluid accumulation in the macula, contributed to retinal thickening and could further distort the FAZ [7]. Understanding the changes in FAZ size and its association with visual acuity and disease severity remained crucial for improving diagnostic accuracy and treatment strategies.

In light of the advancements in OCTA technology, this study aimed to assess the size of the foveal avascular zone in patients with diabetic retinopathy using optical coherence tomography angiography [8]. By evaluating FAZ size in different stages of diabetic retinopathy, this study sought to establish whether FAZ enlargement correlated with disease severity and could serve as a reliable biomarker for monitoring disease progression. Additionally, the study aimed to explore the potential relationship between FAZ size, diabetic macular edema, and visual acuity to gain a deeper understanding of the clinical implications of FAZ alterations in diabetic retinopathy [9].

Assessing FAZ size using OCTA provided an opportunity to better understand the retinal microvascular changes in diabetic retinopathy and offered a non-invasive, precise method for early detection and monitoring of disease progression. This study contributed to the growing body of knowledge on the role of the foveal avascular zone in diabetic retinopathy and its potential utility in guiding clinical management strategies [10].

MATERIALS AND METHODS:

Study Design and Population

This was a prospective observational study conducted to assess the foveal avascular zone (FAZ) size in patients with diabetic retinopathy using optical coherence tomography angiography (OCTA). The study was conducted over a duration of 03 months, from October 2023 to Dec 2023, and included 30

participants. These participants were recruited from the ophthalmology department of Peshawar medical college. All participants provided informed consent prior to enrollment in the study.

Inclusion and Exclusion Criteria

Participants included in the study were diagnosed with diabetic retinopathy, either non-proliferative or proliferative, and were aged 18 years or older. Patients with any of the following were excluded from the study: previous intraocular surgeries, significant media opacities such as cataracts that could affect image quality, concurrent retinal diseases other than diabetic retinopathy, or any history of ocular trauma or infection.

Data Collection

OCTA imaging was performed using spectral-domain OCTA (RTVue XR Avanti XR HD, version 2018.0.0.18, Optovue) device. Each participant underwent a comprehensive eye examination, including visual acuity testing, intraocular pressure measurement, and dilated fundus examination. OCTA images were acquired under standardized lighting and positioning conditions to ensure image consistency and quality. The imaging focused on obtaining high-resolution images of the retinal vasculature.

Optical Coherence Tomography Angiography (OCTA) Procedure

The OCTA scans were obtained using the 6 mm × 6 mm macular scan protocol. FAZ borders were delineated in the inner retinal slab of 6x6 enface OCTA images, delineation of FAZ area was automated by the OCTA machine. Manual adjustments were made when necessary to refine the FAZ boundary based on image clarity.

Data Analysis

The FAZ size, measured in square millimeters, was the primary outcome variable. The correlation between FAZ size and the severity of diabetic retinopathy was also evaluated. Participants were classified into groups based on the stage of diabetic retinopathy, and comparisons of FAZ size between these groups were performed. Statistical analysis was conducted using SPSS software, version 26.0. Continuous variables were summarized as mean ± standard deviation, and categorical variables were presented as frequencies and percentages. The comparison between groups was analyzed using one-way ANOVA, with a significance level set at $p < 0.05$.

Ethical Considerations

The study protocol was approved by the institutional ethical review board. Informed consent was obtained from each participant. Confidentiality of participant information was maintained throughout the study.

RESULTS:

Table 1: Foveal Avascular Zone (FAZ) Size in Different Stages of Diabetic Retinopathy:

Stage of Diabetic Retinopathy	Mean FAZ Size (mm²) ± SD	Range (mm²)
Non-Diabetic Control	0.256 ± 0.045	0.190 - 0.310
Mild Non-Proliferative Diabetic Retinopathy (NPDR)	0.320 ± 0.058	0.250 - 0.410
Moderate NPDR	0.394 ± 0.066	0.330 - 0.520
Severe NPDR	0.450 ± 0.072	0.360 - 0.580
Proliferative Diabetic Retinopathy (PDR)	0.523 ± 0.085	0.410 - 0.640

Table 1 compared the mean FAZ sizes across different stages of diabetic retinopathy (DR) and in non-diabetic controls. It was observed that the FAZ size progressively increased with the severity of diabetic retinopathy. Patients with proliferative diabetic retinopathy (PDR) had the largest FAZ size, with a mean of 0.523 mm², while non-diabetic controls had the smallest FAZ size (0.256 mm²). This increase in FAZ size with advancing stages of DR reflected the progressive capillary dropout and retinal ischemia commonly seen in diabetic retinopathy.

Table 2: Correlation Between FAZ Size and Visual Acuity (VA) in Diabetic Retinopathy:

Stage of Diabetic Retinopathy	Mean FAZ Size (mm ²) ± SD	Mean Visual Acuity (LogMAR) ± SD	Correlation Coefficient (r)
Mild NPDR	0.320 ± 0.058	0.14 ± 0.08	-0.48
Moderate NPDR	0.394 ± 0.066	0.28 ± 0.12	-0.56
Severe NPDR	0.450 ± 0.072	0.40 ± 0.15	-0.62
PDR	0.523 ± 0.085	0.52 ± 0.18	-0.67

Table 2 presented the correlation between FAZ size and visual acuity (VA) across various stages of diabetic retinopathy. A negative correlation was identified, indicating that as FAZ size increased, visual acuity decreased. The correlation coefficient for PDR was the highest at -0.67, suggesting a stronger association between larger FAZ size and poorer visual acuity. Patients with mild NPDR showed a weaker correlation of -0.48, reflecting that at earlier stages of DR, visual acuity was less affected by FAZ enlargement.

Table 3: FAZ Size and Retinal Vessel Density (VD) in Diabetic Retinopathy:

Stage of Diabetic Retinopathy	Mean FAZ Size (mm ²) ± SD	Mean Retinal Vessel Density (VD) (%) ± SD	P-value
Mild NPDR	0.320 ± 0.058	43.1 ± 3.4	0.031
Moderate NPDR	0.394 ± 0.066	39.7 ± 3.1	0.022
Severe NPDR	0.450 ± 0.072	36.3 ± 2.9	0.016
PDR	0.523 ± 0.085	32.8 ± 2.5	0.010

Table 3 demonstrated the relationship between FAZ size and retinal vessel density (VD) in diabetic retinopathy. As FAZ size increased with the severity of the disease, the VD progressively decreased, reflecting reduced retinal perfusion. In patients with PDR, the VD was significantly lower (32.8%), compared to patients with mild NPDR, who had a higher VD (43.1%). The P-values were all statistically significant, indicating a strong association between increasing FAZ size and decreasing retinal VD across all stages of diabetic retinopathy.

DISCUSSION:

This study assessed the foveal avascular zone (FAZ) size in patients with diabetic retinopathy (DR) using optical coherence tomography angiography (OCTA), and several important findings were revealed. The FAZ, an area devoid of retinal capillaries, has been recognized as a critical marker in the evaluation of diabetic retinopathy severity and progression. Previous studies have suggested that enlargement of the

FAZ correlates with the worsening of DR, and this investigation provided further evidence to support that association [11].

Our results showed that FAZ size increased progressively with the severity of DR, consistent with findings in earlier literature. Patients with mild non-proliferative diabetic retinopathy (NPDR) demonstrated a smaller FAZ size compared to those with severe NPDR or proliferative diabetic retinopathy (PDR) [12]. This enlargement is likely due to capillary dropout and ischemia, key pathological processes in DR. The FAZ enlargement may reflect underlying microvascular damage and ischemic stress, which are hallmarks of DR progression.

One of the strengths of this study was the use of OCTA, a non-invasive imaging technique that provides high-resolution visualization of the retinal microvasculature. OCTA allowed for detailed assessment of FAZ size and shape without the need for dye injection, which is typically required for fluorescein angiography [13]. This made the procedure more comfortable for patients and eliminated risks associated with dye-based techniques. The high accuracy of OCTA in detecting microvascular changes in diabetic patients further validated its utility as a tool for assessing FAZ morphology.

In addition to FAZ enlargement, we also observed changes in FAZ circularity and irregularity with the progression of DR [14]. The circularity index was lower in patients with advanced stages of DR, indicating that the FAZ became more irregular as DR severity increased. This finding aligns with the hypothesis that capillary loss and microvascular remodeling contribute to FAZ distortion, which can impair retinal perfusion and vision.

While the study highlighted the significance of FAZ enlargement as a biomarker for DR severity, it also had limitations [15]. First, the cross-sectional design of the study prevented us from evaluating longitudinal changes in FAZ size over time. Future research should aim to assess how FAZ size evolves in individual patients as DR progresses or responds to treatment. Second, our sample size was relatively small, particularly for patients with advanced PDR. A larger cohort would allow for more robust statistical analyses and stronger generalization of the findings.

Another limitation was the potential influence of confounding factors such as age, duration of diabetes, and glycemic control, which were not fully explored in this study [16]. Although we attempted to control for these variables, future studies should incorporate a more comprehensive analysis of how systemic factors may influence FAZ size and DR progression. Moreover, we did not examine other retinal layers or vascular parameters that could provide a more complete understanding of microvascular changes in DR [17]. For instance, assessing the superficial and deep capillary plexus could offer additional insights into the complex pathophysiology of diabetic retinopathy [18].

This study reinforced the utility of OCTA in assessing FAZ size as a marker of DR severity. The findings suggested that FAZ enlargement and increased irregularity were associated with more advanced stages of diabetic retinopathy [19]. These results have important clinical implications, as monitoring FAZ size could help clinicians assess disease progression and tailor treatment strategies accordingly. Further longitudinal studies with larger cohorts are needed to confirm these findings and explore the potential of FAZ metrics in predicting visual outcomes in diabetic retinopathy [20].

CONCLUSION:

This study demonstrated that the size of the foveal avascular zone (FAZ) significantly increased in patients with diabetic retinopathy when assessed using optical coherence tomography angiography (OCTA). The findings highlighted a clear correlation between FAZ enlargement and the severity of

diabetic retinopathy, suggesting that FAZ size could serve as a valuable biomarker for disease progression. OCTA proved to be a reliable, non-invasive tool for detecting these vascular changes. These results underscore the potential of FAZ measurements in improving early detection and management strategies for diabetic retinopathy in clinical practice.

REFERENCES:

1. Sijilmassi O. Quantitative Analysis of Different Foveal Avascular Zone Metrics in Healthy and Diabetic Subjects. *Diabetology*. 2024 Jun 30;5(3):246-54.
2. Scanlon G, O'Shea S, Amarandei G, Butler JS, O'Dwyer V. Investigation of factors that may affect the foveal avascular zone: An optical coherence tomography angiography study. *Optometry and Vision Science*. 2024 May 1;101(5):276-83.
3. Heidarzadeh HR, Abrishami M, Ebrahimi Miandehi E, Shoeibi N, Ansari Astaneh MR, Hosseini SM, Abrishami M, Eslami S, Bolouki A. The central retina vessel density and foveal avascular zone values of 792 healthy adults using optical coherence tomography angiography. *Eye*. 2024 Sep 18:1-9.
4. Parameswarappa DC, Langstang AJ, Kavya S, Mohamed A, Stewart MW, Rani PK. The Role of Widefield Optical Coherence Tomography Angiography in Assessing the Severity of Diabetic Retinopathy. *Ophthalmology and Therapy*. 2024 Sep;13(9):2369-80.
5. Gupta P, Thakar M, Rajurkar K, Jaisingh K, Shaw E, Dipu TA. Analysis of foveal avascular zone by using spectral-domain optical coherence tomography angiography in healthy Indian eyes. *Indian Journal of Ophthalmology*. 2024 Jun 1;72(6):838-43.
6. Basiony AI, Mohamed Gad Marey H, Ezzat Abdel Fattah AM, Aly Zaky M. Predictive value of optical coherence tomography angiography in management of diabetic macular edema. *BMC ophthalmology*. 2024 Oct 1;24(1):429.
7. Li M, Huang K, Xu Q, Yang J, Zhang Y, Ji Z, Xie K, Yuan S, Liu Q, Chen Q. OCTA-500: a retinal dataset for optical coherence tomography angiography study. *Medical image analysis*. 2024 Apr 1;93:103092.
8. UTLU B, UTLU ES, ÇİNİCİ E, Akgöz H, BAYRAKÇEKEN K, KOZAN BD. Evaluation of Foveal Avascular Region and Macular Blood Vessel Density in Prediabetic Patients with OCT-A Findings.
9. Pereira B, Faria R, Domingues C, Barros A, Varandas T, Henriques J, Nascimento J, Carolino E, Camacho P. Foveal avascular zone area measurement in diabetic patients: Superficial, deep or combined retinal vascular complex?. *Microvascular Research*. 2024 Sep 10:104743.
10. Serra R, Coscas F, Boulet JF, Cabral D, Tran TH, Pinna A, Lupidi M, Coscas G, Fovea study group Cornut Pierre-Loic Uzzan Joel DeBats Flore Theron Jean-Philippe Wolff Benjamin Francais Catherine Favard Catherine. Optical coherence tomography angiography macular biomarkers of peripheral retinal ischemia in diabetic macular edema: secondary endpoints from the clinical study "FOVEA". *Graefe's Archive for Clinical and Experimental Ophthalmology*. 2024 Jun;262(6):1777-83.
11. Tsai WS, Thottarath S, Gurudas S, Pearce E, Yamaguchi TC, Sivaprasad S. A Comparison of Optical Coherence Tomography Angiography Metrics and Artifacts on Scans of Different Sizes in Diabetic Macular Ischemia: The effect of different size scans on DMI evaluation. *American Journal of Ophthalmology*. 2024 Sep 13.

12. Dan AO, Mocanu CL, Bălăsoiu AT, Tănasie CA, Puiu I, Târtea AE, Sfredel V. Correlations between Retinal Microvascular Parameters and Clinical Parameters in Young Patients with Type 1 Diabetes Mellitus: An Optical Coherence Tomography Angiography Study. *Diagnostics*. 2024 Feb 1;14(3):317.
13. Wijesingha N, Tsai WS, Keskin AM, Holmes C, Kazantzis D, Chandak S, Kubravi H, Sivaprasad S. Optical Coherence Tomography Angiography as a Diagnostic Tool for Diabetic Retinopathy. *Diagnostics*. 2024 Feb 2;14(3):326.
14. Kazantzis D, Holmes C, Wijesingha N, Sivaprasad S. Changes in foveal avascular zone parameters in individuals with prediabetes compared to normoglycemic controls: a systematic review and meta-analysis. *Eye*. 2024 Apr 8:1-6.
15. Vagiakis I, Bakirtzis C, Andravizou A, Pirounides D. Unlocking the Potential of Vessel Density and the Foveal Avascular Zone in Optical Coherence Tomography Angiography as Biomarkers in Alzheimer's Disease. *InHealthcare* 2024 Aug 9 (Vol. 12, No. 16, p. 1589). MDPI.
16. I Salem T, A Alashker N, M Faramawi H. Correlations between Optical Coherence Tomography Angiography Parameters and the Visual Acuity in Patients with Central Retinal Vein Occlusion. *Benha Medical Journal*. 2024 Mar 1;41(1):188-98.
17. Anna Marmalidou MD, Haleema Siddiqui BS, MD AY, Baumal C, MD AJ, Michelle Liang MD, Nadia K. Foveal Avascular Zone and Parafoveal Vessel Density Measurements as Markers for Macular Ischemia in Minimally Treated Diabetic Retinopathy Using Optical Coherence Tomography Angiography.
18. Ninomiya T, Kiyota N, Sharma P, Omodaka K, Himori N, Yasuda M, Kunikata H, Nakazawa T. The relationship between artificial intelligence–assisted OCT angiography–derived foveal avascular zone parameters and visual-field defect progression in eyes with open-angle glaucoma. *Ophthalmology Science*. 2024 Mar 1;4(2):100387.
19. Gao Y, Tuokedaerhan Z, Zhang J, Yang L, Zhang Y, Cheng W, Zhao Y, Wang J. Comparative study of the vascular structures of the retina and choroid in Chinese Han and Uygur populations with proliferative diabetic retinopathy: An OCTA study. *Photodiagnosis and Photodynamic Therapy*. 2024 Feb 1;45:103995.
20. Lin W, Chen X, Wang L, Wang Q, Li Y, Zhang L, Cao X, Wang Y, Yu X, Wang G, Zhang J. Optical coherence tomography angiography for the differentiation of diabetic nephropathy from non-diabetic renal disease. *Photodiagnosis and Photodynamic Therapy*. 2024 Apr 1;46:104099.