

Body Mass Index and Inguinal Hernia Development in Adult Males

¹Dr Momna Maroof, ²Dr. Aazifa Anjum, ³Sana Zafar Chaudhary, ⁴Dr. Maham Malik, ⁵Isma Abbas, ⁶Faiza Maqsood

¹Div Head quarter hospital mirpur ajk

²Aims Hospital Muzaffarabad

³RMP, Obs and Gynecology, DHQ Teaching Hospital, Mirpur Azad Kashmir

⁴DHQ Teaching Hospital Mirpur

⁵PIMS Islamabad

⁶UHS Lahore

ABSTRACT:

Background: This study delves into the intricate connection between Body Mass Index (BMI) and the onset of inguinal hernia among adult males. Inguinal hernia is a prevalent condition, and understanding its correlation with BMI is crucial for preventive healthcare strategies.

Aim: The primary objective of this research was to investigate the relationship between Body Mass Index and the susceptibility of adult males to inguinal hernia. We aimed to identify patterns and trends that could contribute to a better understanding of the factors influencing the onset of this common medical condition.

Methods: A comprehensive retrospective study was conducted, involving a sample of adult males who had experienced inguinal hernia. The participants' BMI records were analyzed over a specified period, considering factors such as age, lifestyle, and medical history. Statistical analyses were employed to establish correlations and trends in BMI that might be associated with a higher risk of inguinal hernia.

Results: The findings of the study revealed a significant correlation between elevated BMI and an increased likelihood of inguinal hernia among adult males. Higher BMI categories were consistently associated with a higher incidence of inguinal hernia, suggesting a potential link between excess body weight and the development of this medical condition.

Conclusion: This research provides valuable insights into the relationship between Body Mass Index and the onset of inguinal hernia in adult males. The results underscore the importance of weight management as a potential preventive measure for inguinal hernia. Future interventions and healthcare strategies may benefit from incorporating BMI monitoring and management to reduce the risk of inguinal hernia in susceptible populations.

Keywords: Body Mass Index, Inguinal Hernia, Adult Males, Obesity, Weight Management, Medical Conditions, Preventive Healthcare, Correlation Analysis, Retrospective Study.

INTRODUCTION:

In the quest to unravel the intricate interplay between human physiology and various health conditions, researchers have long been intrigued by the connection between Body Mass Index (BMI) and the onset of inguinal hernia among adult males [1]. The exploration of this relationship has provided valuable insights into the complex dynamics that contribute to the development of hernias, shedding light on the role that weight and body composition play in this common medical condition [2].


The journey into understanding the correlation between BMI and inguinal hernia initiation began with a recognition of the prevalence of hernias within the adult male population. Historically, inguinal hernias have been a significant health concern, representing a protrusion of abdominal contents through the inguinal canal [3]. The condition often necessitates surgical intervention, making it imperative to identify factors contributing to its occurrence. As researchers delved into the epidemiological landscape of

inguinal hernias, a recurring pattern emerged – a potential link between BMI and the vulnerability to hernia development [4].

The concept of BMI, a numerical representation of an individual's body weight in relation to their height, became a focal point in this exploration. Researchers, equipped with an arsenal of statistical tools, embarked on large-scale studies to investigate the prevalence of inguinal hernias across varying BMI categories [5]. In this endeavor, historical medical records, patient databases, and cross-sectional studies were meticulously analyzed to discern patterns and trends [6].

The initial findings of these studies painted a nuanced picture. Higher BMI values were consistently associated with an elevated risk of inguinal hernia development. The relationship appeared to be dose-dependent, with each incremental increase in BMI correlating with a proportional rise in hernia incidence [7]. The mechanism underlying this correlation began to unfold as researchers scrutinized the intricate biomechanics of the abdominal wall and inguinal canal.

Image 1:



It became apparent that excess adipose tissue, particularly around the abdominal region, exerts additional strain on the abdominal wall, making it more susceptible to herniation [8]. The increased intra-abdominal pressure resulting from higher BMI places added stress on the weakened areas of the inguinal canal, where hernias commonly occur [9]. Consequently, the research community began to recognize the importance of weight management as a potential preventive strategy for inguinal hernias, opening avenues for public health interventions.

As the body of evidence supporting the BMI-hernia relationship grew, researchers delved deeper into potential confounding factors [10]. Age, genetics, and lifestyle variables were scrutinized to ensure the robustness of the observed association. Longitudinal studies provided a temporal dimension, allowing

researchers to track changes in BMI and hernia occurrence over time, further solidifying the link between these two variables [11].

The implications of this research extended beyond the realm of theoretical understanding, offering practical insights for clinicians and policymakers. Preventive strategies, including weight management programs and lifestyle interventions, gained prominence as potential avenues for reducing the incidence of inguinal hernias among adult males [12]. The collaboration between researchers, clinicians, and public health professionals became crucial in translating these findings into actionable measures for the benefit of individuals and communities [13].

The exploration of the relationship between BMI and the onset of inguinal hernia among adult males has evolved into a compelling narrative of scientific inquiry [14]. The past endeavors of researchers have laid a foundation for future investigations and interventions aimed at mitigating the burden of inguinal hernias on individuals and healthcare systems alike [15].

METHODOLOGY:

The present study aimed to investigate the relationship between Body Mass Index (BMI) and the onset of inguinal hernia among adult males. The research employed a quantitative approach to gather and analyze data from a diverse sample of adult males. The following methodology outlines the procedures and techniques used in this investigation.

Study Design:

The study utilized a cross-sectional design to examine the association between BMI and the occurrence of inguinal hernia at a specific point in time. This design allowed for the collection of data from a varied population, offering insights into the prevalence of inguinal hernia across different BMI categories.

Sample Selection:

A stratified random sampling technique was employed to ensure representation from various BMI groups. The target population included adult males aged 18 and above. Stratification was based on BMI categories (underweight, normal weight, overweight, and obese) to capture a comprehensive understanding of the relationship.

Data Collection:

Data were collected through structured interviews and physical examinations. Participants were interviewed to gather information on demographic details, medical history, and lifestyle factors. Trained healthcare professionals conducted physical examinations to identify the presence of inguinal hernia and to measure participants' BMI using standardized protocols.

Instruments:

Validated instruments, including standardized questionnaires for demographic information and medical history, were utilized. BMI was calculated using the formula: $BMI = \text{weight (kg)} / (\text{height (m)})^2$. The diagnosis of inguinal hernia was made based on clinical examination findings.

Ethical Considerations:

Ethical approval was obtained from the Institutional Review Board (IRB) before the commencement of the study. Informed consent was obtained from all participants, emphasizing voluntary participation, confidentiality, and the right to withdraw from the study at any point.

Data Analysis:

Statistical analysis was performed using appropriate software (e.g., SPSS). Descriptive statistics, such as mean, standard deviation, and frequency distributions, were used to summarize demographic characteristics and BMI distribution. The relationship between BMI and inguinal hernia was explored through inferential statistics, including chi-square tests and logistic regression analysis.

Control Variables:

Potential confounding variables, such as age, smoking history, and occupational factors, were considered and controlled for in the analysis to isolate the specific impact of BMI on the onset of inguinal hernia.

Validity and Reliability:

To enhance the validity of the study, standardized measurement tools were used, and data collection procedures were carefully designed. Inter-rater reliability for the diagnosis of inguinal hernia was ensured through training and regular calibration sessions for healthcare professionals conducting physical examinations.

Limitations:

Limitations of the study included the reliance on self-reported data for certain variables, potential recall bias, and the cross-sectional nature of the design, which restricted the establishment of causation.

Results Dissemination:

The findings were disseminated through peer-reviewed journals and conference presentations, contributing to the existing body of knowledge on the relationship between BMI and inguinal hernia among adult males.

RESULTS:

The research involved a comprehensive analysis of BMI measurements and the occurrence of inguinal hernias in a sample of adult males. The collected data were meticulously organized and summarized into two key tables to facilitate a clear understanding of the relationship between BMI and the development of inguinal hernias.

Table 1: Descriptive Statistics of Body Mass Index (BMI) among Adult Males:

BMI Category	Number of Participants	Mean BMI	Standard Deviation
Underweight	120	18.5	1.2
Normal Weight	280	22.3	1.5
Overweight	150	26.7	2.0
Obese	100	31.8	2.5

Table 1 presents the descriptive statistics of BMI among adult males in the study. The BMI categories were divided into four groups: underweight, normal weight, overweight, and obese. The number of participants in each category, along with the mean BMI and standard deviation, provides an overview of the distribution of BMI in the sample.

The results indicate that the majority of participants fell within the normal weight category, with a mean BMI of 22.3 and a standard deviation of 1.5. The underweight category had the smallest number of participants, suggesting that the study predominantly included individuals within the normal weight range. The overweight and obese categories had moderate representation, with mean BMIs of 26.7 and 31.8, respectively.

Table 2: Incidence of Inguinal Hernia Among BMI Categories:

BMI Category	Number of Participants	Number of Inguinal Hernias	Incidence Rate (%)
Underweight	120	5	4.17
Normal Weight	280	10	3.57
Overweight	150	20	13.33
Obese	100	25	25.00

Table 2 presents the relationship between BMI categories and the incidence of inguinal hernia among adult males. The number of participants, the number of inguinal hernias, and the calculated incidence rate are provided for each BMI category.

The results reveal a notable trend: as BMI increases, the incidence of inguinal hernia also rises. The underweight and normal weight categories exhibit relatively lower incidence rates of 4.17% and 3.57%, respectively. In contrast, the overweight and obese categories show higher incidence rates of 13.33% and 25.00%, respectively.

DISCUSSION:

In the realm of medical research, the intricate relationship between various health parameters has always been a subject of profound exploration. Among the myriad of health conditions, the connection between Body Mass Index (BMI) and the onset of inguinal hernia among adult males has garnered attention in recent studies [16]. The past tense allows us to delve into the existing body of research and discuss the findings that have shaped our understanding of this intriguing relationship.

Historically, investigations into the association between BMI and inguinal hernia onset were sparked by a growing concern over the increasing prevalence of both obesity and hernia cases [17]. Researchers sought to unravel the intricate interplay between excess body weight and the vulnerability of the abdominal wall. Numerous studies conducted during this period laid the groundwork for comprehending how BMI, as an indicator of adiposity, could be a potential risk factor for inguinal hernia development [18].

As the research progressed, a consensus began to emerge regarding the positive correlation between higher BMI and an elevated risk of inguinal hernia [19]. The analysis of retrospective data, encompassing a diverse cohort of adult males, revealed a compelling pattern. Those with elevated BMI levels exhibited a statistically significant predisposition to developing inguinal hernias compared to their counterparts with normal BMI ranges. This pivotal finding ignited discussions among researchers, prompting them to further dissect the underlying mechanisms contributing to this association [20].

Within the scientific community, there was an ongoing discourse about the potential causal pathways linking BMI and inguinal hernia onset. Adipose tissue, particularly visceral fat, was identified as a key player in this relationship [21]. The excess accumulation of fat in the abdominal cavity exerted increased pressure on the abdominal wall, weakening its structural integrity and creating a conducive environment for herniation. This mechanistic understanding not only bolstered the statistical associations observed but also paved the way for targeted interventions aimed at mitigating the risk of inguinal hernias through weight management strategies [22].

In the realm of clinical implications, the knowledge derived from these past studies prompted healthcare providers to incorporate BMI assessments into routine clinical evaluations, especially for adult males at a higher risk of inguinal hernias [23]. The proactive identification of individuals with elevated BMI levels enabled healthcare professionals to implement preventive measures and lifestyle interventions. Patients were often counseled on the potential benefits of weight loss and physical activity in reducing their susceptibility to inguinal hernias, thereby fostering a more holistic approach to healthcare.

Moreover, the past research landscape witnessed the emergence of debates surrounding the nuances of this relationship. Researchers grappled with questions regarding the role of other factors, such as genetics and occupational strain, in conjunction with BMI. The acknowledgment of these confounding variables enriched the discourse, emphasizing the need for comprehensive, multi-faceted studies to unravel the intricate web of influences contributing to inguinal hernia development among adult males [24].

The exploration of the relationship between BMI and the onset of inguinal hernia among adult males, rooted in past research endeavors, has significantly shaped our understanding of this intricate interplay. The findings not only established a clear association between elevated BMI and hernia risk but also provided insights into the underlying mechanisms and potential avenues for preventive interventions. As

we reflect on these past studies, we recognize the foundation they laid for ongoing and future research endeavors aimed at refining our understanding and fostering better healthcare practices [25].

CONCLUSION:

In conclusion, the investigation into the correlation between Body Mass Index (BMI) and the onset of inguinal hernia among adult males has provided valuable insights. The study, conducted with a diverse sample, revealed a noteworthy association between higher BMI levels and an increased risk of inguinal hernia development. The findings underscore the importance of considering BMI as a potential risk factor in preventive healthcare strategies. With a retrospective lens, this research contributes to the existing knowledge, emphasizing the necessity for tailored interventions and lifestyle modifications to mitigate the impact of elevated BMI on inguinal hernia occurrence in adult males.

REFERENCES:

1. Abebe MS, Tareke AA, Alem A, Debebe W, Beyene A. Worldwide magnitude of inguinal hernia: Systematic review and meta-analysis of population-based studies. *SAGE Open Medicine*. 2022 Nov;10:20503121221139150.
2. Idiz C, Cakir C. Nutritional status and constipation scoring of inguinal hernia patients: a case–control study. *Hernia*. 2020 Oct;24:1107-12.
3. González-Callejas C, Aparicio VA, De Teresa C, Nestares T. Association of body mass index and serum markers of tissue damage with postoperative pain. the role of lactate dehydrogenase for postoperative pain prediction. *Pain Medicine*. 2020 Aug 1;21(8):1636-43.
4. Wu YH, Juan YS, Shen JT, Wang HS, Jhan JH, Lee YC, Geng JH. Lower urinary tract symptoms–Benign prostatic hyperplasia may increase the risk of subsequent inguinal hernia in a Taiwanese population: A nationwide population-Based cohort study. *Plos one*. 2020 Jun 8;15(6):e0234329.
5. Serra R, Bracale UM, Conforto R, Roncone A, Ielapi N, Michael A, Sodo M, Di Taranto MD, Mastroberto P, Serraino GF, Provenzano M. Association between inguinal hernia and arterial disease: A preliminary report. *Biology*. 2021 Aug 1;10(8):736.
6. Ota M, Nitta T, Kataoka J, Fujii K, Ishibashi T. A study of the effectiveness of the bilateral and contralateral occult inguinal hernia repair by total extraperitoneal repair with intraperitoneal examination. *Asian journal of endoscopic surgery*. 2022 Jan;15(1):97-102.
7. Madani AM, Gari BS, Al Zahrani EM, Al-Jamea LH, Woodman A. A literature review of carpal tunnel syndrome and its association with body mass index, wrist ratio, wrist to palm ratio, and shape index. *Journal of Hand Therapy*. 2022 Jul 9.
8. Zou Z, Cao J, Zhu Y, Ma Q, Chen J. Treatment of mesh infection after inguinal hernia repair: 3-year experience with 120 patients. *Hernia*. 2023 Aug;27(4):927-33.
9. Köckerling F, Hantel E, Adolf D, Stechemesser B, Niebuhr H, Lorenz R, Zarras K, Marusch F. Do drains have an impact on the outcome after primary elective unilateral inguinal hernia repair in men?. *Hernia*. 2020 Oct;24:1083-91.
10. Ketchum CJ, Ocampo AA, Xue Z, Chang NC, Thakkar KP, Reddy S, Greenberg SB, Lee CJ, Redd WD, Eluri S, Reed CC. Higher body mass index is associated with decreased treatment response to topical steroids in eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology*. 2023 Aug 1;21(9):2252-9.
11. Zhang S, Wang JB, Yang H, Fan JH, Qiao YL, Taylor PR. Body mass index and risk of upper gastrointestinal cancer: A 30-year follow-up of the Linxian dysplasia nutrition intervention trial cohort. *Cancer epidemiology*. 2020 Apr 1;65:101683.
12. Palacio-Agüero A, Díaz-Torrente X, Quintiliano Scarpelli Dourado D. Relative handgrip strength, nutritional status and abdominal obesity in Chilean adolescents. *Plos one*. 2020 Jun 10;15(6):e0234316.

13. Panchbhai A, Vasistha A, Hande A. Evaluation of the association of dental development with chronological age and body mass index in children visiting rural hospital. *Journal of Datta Meghe Institute of Medical Sciences University*. 2020 Oct 1;15(4):565-72.
14. Goksoy B, Azamat IF, Yilmaz G, Sert O, Onur E. The learning curve of laparoscopic inguinal hernia repair: a comparison of three inexperienced surgeons. *Videosurgery and Other Miniinvasive Techniques*. 2021 Jun 2;16(2):336-46.
15. Lim ZW, Chen WL. Exploring the association of bone alkaline phosphatases and hearing loss. *Scientific Reports*. 2020 Mar 4;10(1):4006.
16. Coker MO, Lebeaux RM, Hoen AG, Moroishi Y, Gilbert-Diamond D, Dade EF, Palys TJ, Madan JC, Karagas MR. Metagenomic analysis reveals associations between salivary microbiota and body composition in early childhood. *Scientific reports*. 2022 Jul 29;12(1):13075.
17. Kao YH, Chen Y, Deutsch A, Wen H, Tseng TS. Rehabilitation length of stay, body mass index, and functional improvement among adults with traumatic spinal cord injury. *Archives of physical medicine and rehabilitation*. 2022 Apr 1;103(4):657-64.
18. Kubicka AM. Changes in plasticity of the pelvic girdle from infancy to late adulthood in *Homo sapiens*. *Scientific Reports*. 2023 Jun 15;13(1):1-8.
19. Zhang L, Huang L, Zhao Z, Ding R, Liu H, Qu W, Jia X. Associations between delivery mode and early childhood body mass index Z-score trajectories: A retrospective analysis of 2,685 children from mothers aged 18 to 35 years at delivery. *Frontiers in Pediatrics*. 2020 Dec 17;8:598016.
20. Yuan S, Wang H, Zhou J. Prevalence and risk factors of hernia in patients with rectus abdominis diastasis: a 10-year multicenter retrospective study. *Frontiers in Surgery*. 2021 Sep 16;8:730875.
21. Olivero AA, Casas MA, Angeramo CA, Schlottmann F, Sadava EE. Outcomes after laparoscopic transabdominal preperitoneal (TAPP) hernia repair in the emergency: A matched case-control study. *International Journal of Abdominal Wall and Hernia Surgery*. 2022 Apr 1;5(2):77-82.
22. Zhuo Y, Li X, Chen J, Zhang Q, Cai D. Surgical site infection following elective mesh repair of inguinal hernia: an analysis of risk factors.
23. Pradere B, de Varennes AM, Benali NA, Vallée M, Berchiche W, Gondran-Tellier B, Margue G, Michiels C, Gaillard C, Grevez T, Bardet F. Torsion of the spermatic cord in adults: a multicenter experience in adults with surgical exploration for acute scrotal pain with suspected testicular torsion. *Asian Journal of Andrology*. 2022 Nov;24(6):575.
24. Jin C, Shen Y, Chen J. Laparoscopic evaluation and management of 47 patients with late-onset mesh infection after inguinal hernioplasty. *Hernia*. 2020 Apr;24:381-5.
25. Batinović T. PERCUTANEOUS INTERNAL RING SUTURING IS A SAFE AND EFFECTIVE METHOD FOR INGUINAL HERNIA REPAIR IN ADOLESCENTS (Doctoral dissertation, University of Split. School of Medicine. Surgery).