

THE FREQUENCY AND CHARACTERISTICS OF POSTSURGICAL PROBLEMS IN INDIVIDUALS WHO HAD GOT NEUROSURGICAL OPERATIONS

¹Muhammad Musa, ²Farnaz Ahsan, ³Dr Muhammad Salman Tahir, ⁴Sadam Hussain, ⁵Fayiz F El- Shamy, ⁶Samreen KishwarAbbasi

¹Services Institute of Medical Sciences / Services Hospital, Lahore

²DHQ Teaching Hospital Dera Ghazi Khan

³MO, Hameed Clinic Nishtar Road Multan

⁴Medical Officer, Three gorges Company Medical and Surgical Complex Karot, Rwp,

⁵Department of Physical Therapy for Women Health, Kafrelsheikh University Egypt,

⁶WMO, Medical, Ayub Teaching Hospital Abbottabd

ABSTRACT:

Aim: The main aim of our current research was to assess occurrence and prevalence of postsurgical problems in individuals who had had neurosurgical operations.

Methods: Throughout a four-month timeframe, most patients taking neurosurgery remained monitored surgically in post anesthesia care unit or critical care unit for up to four hours. The investigators noted relevant problems and clinical records on a standardized form. Respiratory, cardiovascular, nausea and vomiting, hypothermia, and other problems have been identified. Age, gender, ASA status, kind of surgery, elective or emergency surgery, in addition postoperative placement were all examined as lifestyle factors for the incidence of problems.

Results: Six hundred patients treated studied monitored, although the trachea remained deeply sedated in 57 of them after the four-hour study period, hence they were eliminated from the postpartum effects assessment. At least one situation happened in 56.6 percent of the remaining 437 individuals. Respiratory issues occurred in 3.9 percent of cases, airway damage in 5.6 percent of cases, cardiovascular complications in 7.8 percent of cases, neurological abnormalities in 6.8 percent of cases, and nausea and/or vomiting in 39 percent of cases. The most patients had problems throughout spine (68%) and vascular (68%) surgery, comparing to tumor (48%) and other (47%) surgery, P 0.06. Other lifestyle factors for nausea and vomiting were being over 70 years old (P 0.01) and having elective spine also vascular surgery (P 0.002).

Conclusion: In patients presenting, here had been very significant rate of early postoperative problems. The most prevalent issue included nausea and vomiting, that was more prevalent among young patients receiving elective spine surgery.

Keywords: Occurrence and prevalence of postsurgical problems, neurosurgical operations.

7997

DOI Number: 10.14704/nq.2022.20.8.NQ44825

NeuroQuantology 2022; 20(8): 7997-8001

INTRODUCTION:

Patients having a greater risk of negative outcomes during the early postoperative phase. Numerous studies and surveys have indicated an incidence of acute perioperative problems ranging from 4% to 33% in the general surgical population [1]. The most common side effects are respiratory, cardiovascular, and nausea and vomiting. The scope and severity of these problems in the neurosurgical population are not fully established [2]. The goal of this research was to determine the frequency and patterns of problems that arise in immediate postoperative period in adult cases receiving neurosurgical operations. In our study, 6.5 percent of cases had an intraoperative issue that remained identical to postoperative condition. Unfortunately, our data on intraoperative occurrences remained restricted since we only knew what an anesthetist had recorded on anesthetic record [3]. Neurological problems happened in 6.8 percent of patients, which is not surprising given the patient population. Previous studies did not include a patient population receiving neurosurgical operations since the majority among those individuals would also have been hospitalized to the intensive care unit and therefore eliminated from their analyses. Previous research has looked into a variety of risk factor for development of postoperative problems [4]. This included the patient's age, gender, location of surgery, length of operation, ASA status, anesthetic method, and whether the treatment was emergency or elective. In our investigation, being 71 years old was the dangerous aspect for nausea and vomiting, but not for other problems. These are in contrast to earlier study

that found that being above the age of 60 was the dangerous aspect for respiratory and cardiovascular problems [5].

METHODOLOGY:

The institutional independent commission gave their consent from February 2021 to January 2022, all sequential patients receiving a

neur
osurg
ical
oper
ation
are
inclu
ded
in the.
Funct
ional
electi
ve
surge
ries
such
as
chro
nic
pain
and
mobil
ity
probl
ems
were
not
cover
ed.
The
durat
ion of
the
perio
perat
ive
inves
tigati

on was confined to first five hours in either post anesthesia care unit or the critical care unit. Here was not any effort to modify or affect routine patient care practice. A computational chemist and/or the scientists utilized a predefined, standardized form to gather all of the material. Demographic information, analysis, surgical technique, weather process was elective or emergency, and patient's diagnostic examination, including neurological condition, were included with the material. The nurse assessed and documented all vital signs and neurological assessments once the patient arrived in PACU or ICU. There were no early access impairments discovered. This data, as well as any unusual occurrences or problems that happened throughout first four hours in PACU or ICU, was recorded on the designated form. The intraoperative course too was examined, and any unusual occurrences noted on the anesthesia chart became reported. Table I lists the postoperative complications that were evaluated in this review. This research did not include a careful evaluation of postoperative discomfort. The time of extubating was noted if the person arrives with both the trachea heavily sedated. Participants whose lungs were electively ventilated postoperatively and one whose trachea remained hospitalized for the whole four-hour period were omitted from the research methodology for problems. Tukey-type multiple comparison testing was used to examine the gender rates in patients with difficulties and the percentage of difficulties in each group. Chi-square analysis remained performed to evaluate the connection between the incidence of complications and each of following parameters: age, gender, ASA status, elective or emergency surgery, kind of surgery, and postpartum placement in order to

recognize possible risks for the frequency of

7998

problems. SPSS version 7.1.3 was used for the descriptive statistics.

RESULTS:

This research enrolled 481 participants. Additional 27 patients' data sheets appeared insufficient and were not considered. The method of anesthesia and postoperative insertion were performed in accordance having standard practice. The surgeon made the decision whether to use anesthesia or conscious sedation having local anesthetic based on the type of treatment being done. Altogether awake craniotomies for tumors were performed at the surgeon's request for those patient characteristics. All following surgeries were carried out under local anesthetic. The preoperative placement of patients was done in accordance with established hospital policy. The cases were transferred immediately to the ICU following craniotomy under general anesthesia and all vascular operations. Patients recovered in the PACU for four hours after an awake craniotomy before being moved to the ward. Patients who underwent stereotactic biopsy being cared for in the intensive care unit. All additional operations were dictated by the child's perioperative status. The participants included split into four primary surgical categories for assessment: brain tumor (135), spinal surgery (containing spinal tumors) (168), vascular surgery (57), and other (134) (Table II). The fourth category includes the wide range of operations, involving CSF draining (39), peripheral nerve (29), hematoma burr hole (19), ventriculostomy (18), Omiya reservoir (13), and others (19). In the postnatal period, 48 percent of patients recovered in the PACU. Three

individuals were voluntarily admitted to ICU for awake craniotomies. Postoperative period, 32 percent of spine and 47 percent of the other group were transported immediately to ICU. In total, 101 operations (21%) were classed as emergency (Table II). The tracheas were intubated and lungs evacuated for length of the research in 58 of the total (489) patients, excluding them from the definition of the problem. 37 (66%) of them were safety protocols. The severe postoperative status of 37 patients was the explanation for the patients who require intubation and ventilation (brain tumor 6, spine 4, vascular 4, other 26). Since the procedure was protracted, complicated, or unstable, the trachea was electively left anaesthetized in 25 patients. The study of the issues listed in Table I that happened in the 439 individuals analyzed revealed that 56.6 percent of the patients had at least one complication. In total, 101 operations (21%) were classed as emergency (Table II). The tracheas were intubated and the lungs evacuated for the length of the research in 58 of the total (489) patients, excluding them from the definition of the problem. 37 (66%) of them were safety protocols. The severe postoperative status of 37 patients was the explanation for the patients who require intubation and ventilation (brain tumor 6, spine 4, vascular 4, other 26). Since the procedure was protracted, complicated, or unstable, the trachea was electively left anaesthetized in 25 patients. The study of the issues listed in Table I that happened in the 439 individuals analyzed revealed that 56.6 percent of cases had at least one problem.

7999

Table 1:


Group	n	Gender	Age	Emergency	Ventilated
Tumor n (134)	22	14/10	54+20	2	0
	83	39/46	49+19	11	11
	25	12/19	58+17	4	13
Spine (n=175)	63	42/22	51+18	9	3
	108	58/50	15+13	7	3
Others	16	9/10	68+64	1	1

(n=132)	14	15/13	42+22	8	6
---------	----	-------	-------	---	---

Table 2:

Group n	Spine	Tumor	Vascular	Other	Total 450
1 Complications	69 (45%)	46 (37%)	26 (52%)	43 (41%)	183 (42.8%)
2 Complications	13 (19.5%)	12 (9.8 %)	7 (13.5%)	4 (4.98%)	51 (13.5%)
3 Complications	5 (3.6%)	3 (2.5%)	5 (3.6%)	2 (5%)	8 (2.7%)
4 Complications	103 (66%)	65 (48%)	35 (67%)	48 (46%)	238(55.8%)
Total Complication	145 (91%)	75 (60%)	41 (84%)	49 (51%)	302 (71%)

Image 1:



8000

DISCUSSION:

They discovered that the overall prevalence of initial postoperative problems in cases receiving neurosurgical operations was greater than previously stated for general surgical populace in our research. Problems were recorded in 4.3

percent and 6.8 percent of 117,728 anesthetics during two separate time periods in a study of 116,725 anesthetics [6]. The rise in second period was attributable to improvements in anesthetic practice, including growing usage of balanced anesthesia also surveillance. The higher overall prevalence in our study's

neurosurgical group might well be attributed to our limited sample size, data capture technique, high occurrence of nausea and vomiting, or because this proportion of people is more prone to problems in early postoperative phase [7]. The most prevalent problem in our study was nausea and vomiting, particularly happened more often than in prior research. The occurrence of respiratory and cardiovascular problems, on the other hand, was not significant. Our trial began four hours following our admission in the PACU or ICU. This period is selected to correspond with our

repetition [8]. Individuals who have significant operations but do not require ICU, including awake craniotomy and cervical spine surgery, are typically watched in PACU for five hours before being moved to ward. Respiratory issues are concerning in somewhat patients, but they are more so in the neurosurgical doctor [9]. Previous studies indicate a prevalence ranging from 1.4 to 7.8 percent. In our study of 18,067 individuals, we discovered that the incidence of early respiratory problems was 3.96 percent. They discovered that the occurrence of problems was connected to the state of awareness at arrival times, regardless of ASA or age. Hypoventilation, airway blockage, and persistent numbness were the most prevalent issues. A major respiratory episode occurred in 2.4 percent of the 28,186 individuals who underwent a general anesthesia, according to researchers [10].

CONCLUSION:

In conclusion, neurosurgical individuals are susceptible to postoperative problems. The rate of respiratory and cardiovascular actions in PACU or ICU is comparable to that of the overall population. The most common issues in this category are nausea and vomiting, particularly in younger patients who were treated spine surgery.

REFERENCES:

1. B.K. Lal, N.K. Prasad, B.R. Englum, *et al.* Perioperative complications in patients with SARS-CoV-2 infection compared to those without infection: a nationwide propensity-matched analysis
2. Am J Surg (S0002-9610-1) (2020), [10.1016/j.amjsurg.2020.12.024](https://doi.org/10.1016/j.amjsurg.2020.12.024)In Press F. Doglietto, M. Vezzoli, F. Gheza, *et al.* Factors associated with surgical mortality and complications among

patients with and without coronavirus disease 2019 (COVID-19) in Italy *JAMA Surg* (2020), [10.1001/jamasurg.2020.2713](https://doi.org/10.1001/jamasurg.2020.2713)

3. S. Lei, F. Jiang, W. Su, *et al.* Clinical characteristics and outcomes of patients undergoing surgeries during the incubation period of COVID-19

i
n
f
e
c
t
i
o
n
E
C
l
i
n
i
c
a
l
M
e
d
i
c
i
n
e,
2
1
(
2
0
2
0)
,
p.
1
0
0
3
3
1,
1

[0.1016/j.eclinm.2020.100331](https://doi.org/10.1016/j.eclinm.2020.100331)

Article e38, [10.1056/nejmp2015897](https://doi.org/10.1056/nejmp2015897)

4. J.C. Glasbey, A. Bhangu Elective cancer surgery in COVID-19–free surgical pathways during the SARS-CoV-2 pandemic: an international, multicenter, comparative cohort study
J ClinOncol, 20 (October 2020), Article 01933, [10.1200/jco.20.01933](https://doi.org/10.1200/jco.20.01933)
5. COVID Surgery Collaborative. Preoperative nasopharyngeal swab testing and postoperative pulmonary complications in patients undergoing elective surgery during the SARS-CoV-2 pandemic. Br J Surg. doi:10.1093/bjs/znaa051.
6. American College of Surgeons COVID-19: guidance for triage of non-emergent surgical procedure
<https://www.facs.org/covid-19/clinical-guidance/triage>, Accessed 28th Sep 2020
7. Royal College of Surgeons Covid-19: good practice for surgeons and surgical teams
<https://www.rcseng.ac.uk/standards-and-research/standards-and-guidance/good-practice-guides/coronavirus/>, Accessed 28th Sep 2020
8. B.W. Buchan, J.S. Hoff, C.G. Gmehlin, *et al.* Distribution of SARS-CoV-2 PCR cycle threshold values provide practical insight into overall and target-Specific sensitivity among symptomatic patients Am J ClinPathol, 154 (4) (2020), pp. 479-485, [10.1093/AJCP/AQAA133](https://doi.org/10.1093/AJCP/AQAA133)
9. R. Arnaout, R. Lee, G.R. Lee, *et al.* SARS-CoV2 testing: the limit of detection matters bioRxivPreprServBiol (2020), pp. 617-667, [10.1101/2020.06.02.131144](https://doi.org/10.1101/2020.06.02.131144)
10. S. Woloshin, N. Patel, A.S. Kesselheim False negative tests for SARS-CoV-2 infection — challenges and implications N Engl J Med, 383 (6) (2020),

8001